

Michael Karas
Publications 2006-2011

- Aivaliotis M, Haase W, **Karas M**, Tsiotis G (2006) Proteomic analysis of chlorosome-depleted membrane of the green sulfur bacterium *Chlorobium tepidum*. *Proteomics* 6:217-232
- Aivaliotis M, **Karas M**, Tsiotis G (2006) High throughput two-dimensional blue-native electrophoresis: a tool for functional proteomics of cytoplasmatic protein complexes from *Chlorobium tepidum*. *Photosynth Res* 88:143-157
- Aivaliotis M, **Karas M**, Tsiotis G (2007) An alternative strategy for the membrane proteome analysis of the green sulfur bacterium *Chlorobium tepidum* using blue native PAGE and 2-D PAGE on purified membranes. *J Proteome Res* 6:1048-1058
- Andre M, **Karas M** (2007) Investigation of sample-purification procedures for MALDI-based proteomic studies. *Anal Bioanal Chem* 389:1047-1053
- Arrey TN, Rietschel B, Papatotiriou DG, Bornemann S, Baeumlisberger D, **Karas M**, Meyer B (2010) Approaching the complexity of elastase-digested membrane proteomes using Off-Gel IEF/nLC-MALDI-MS/MS. *Anal Chem* 82:2145-2149
- Baeumlisberger D, Arrey TN, Rietschel B, Rohmer M, Papatotiriou DG, Mueller B, Beckhaus T, **Karas M** (2010) Labeling elastase digests with TMT: Informational gain by identification of poorly detectable peptides with MALDI-TOF/TOF mass spectrometry. *Proteomics* 10:3905-3909
- Baeumlisberger D, Rohmer M, Arrey TN, Mueller BF, Beckhaus T, Bahr U, Barka G, **Karas M** (2011) Simple Dual-Spotting Procedure Enhances nLC-MALDI MS/MS Analysis of Digests with Less Specific Enzymes. *J Proteome Res* 10:2889-2894
- Bahr U, Aygun H, **Karas M** (2008) Detection and relative quantification of siRNA double strands by MALDI mass spectrometry. *Anal Chem* 80:6280-6285
- Bahr U, Aygun H, **Karas M** (2009) Sequencing of single and double stranded RNA oligonucleotides by acid hydrolysis and MALDI mass spectrometry. *Anal Chem* 81:3173-3179
- Benedikt A, Baltruschat S, Scholz B, Bursen A, Arrey TN, Meyer B, Varagnolo L, Muller AM, **Karas M**, Dingermann T, Marschalek R (2011) The leukemogenic AF4-MLL fusion protein causes P-TEFb kinase activation and altered epigenetic signatures. *Leukemia* 25:135-144
- Bluem R, Schmidt E, Corvey C, **Karas M**, Schlicksupp A, Kirsch J, Kuhse J (2007) Components of the translational machinery are associated with juvenile glycine receptors and are redistributed to cytoskeleton upon aging and synaptic activity. *J Biol Chem* 282:37783-37793
- Bornemann S, Rietschel B, Baltruschat S, **Karas M**, Meyer B (2010) A novel polyacrylamide gel system for proteomic use offering controllable pore expansion by crosslinker cleavage. *Electrophoresis* 31:585-92
- Burré J, Beckhaus T, Corvey C, **Karas M**, Zimmermann H, Volkandt W (2006) Synaptic vesicle proteins under conditions of rest and activation: Analysis by 2-D difference gel electrophoresis. *Electrophoresis* 27:3488-3496
- Burré J, Beckhaus T, Schagger H, Corvey C, Hofmann S, **Karas M**, Zimmermann H, Volkandt W (2006) Analysis of the synaptic vesicle proteome using three gel-based protein separation techniques. *Proteomics* 6:6250-6262
- Ejsing CS, Moehring T, Bahr U, Duchoslav E, **Karas M**, Simons K, Shevchenko A (2006) Collision-induced dissociation pathways of yeast sphingolipids and their molecular profiling in total lipid extracts: a study by quadrupole TOF and linear ion trap-orbitrap mass spectrometry. *J Mass Spectrom* 41:372-389

Flender C, Leonhard P, Wolf C, Fritzsche M, **Karas M** (2010) Analysis of boronic acids by nano liquid chromatography-direct electron ionization mass spectrometry. *Anal Chem* 82:4194-200

Fuchs SW, Jaskolla TW, Bochmann S, Kotter P, Wichelhaus T, **Karas M**, Stein T, Entian KD (2011) Entianin, a novel subtilin-like lantibiotic from bacillus subtilis subsp spizizenii DSM 15029(T) with high antimicrobial activity. *Appl Environ Microb* 77:1698-1707

Galkin A, Meyer B, Wittig I, **Karas M**, Schägger H, Vinogradov A, Brandt U (2008) Identification of the mitochondrial ND3 subunit as a structural component involved in the active/deactive enzyme transition of respiratory complex I. *J Biol Chem* 283:20907-20913

Hartmann K, Corvey C, Skerka C, Kirschfink M, **Karas M**, Brade V, Miller JC, Stevenson B, Wallich R, Zipfel PF, Kraiczky P (2006) Functional characterization of BbCRASP-2, a distinct outer membrane protein of Borrelia burgdorferi that binds host complement regulators factor H and FHL-1. *Mol Microbiol* 61:1220-1236

Hartshorne RS, Kern M, Meyer B, Clarke TA, **Karas M**, Richardson DJ, Simon J (2007) A dedicated haem lyase is required for the maturation of a novel bacterial cytochrome c with unconventional covalent haem binding. *Mol Microbiol* 64:1049-1060

Herget M, Oancea G, Schrodt S, **Karas M**, Tampé R, Abele R (2007) Mechanism of substrate sensing and signal transmission within an ABC transporter - Use of a trojan horse strategy. *J Biol Chem* 282:3871-3880

Hoang T, Roth U, Kowalewski K, Belisle C, Steinert K, **Karas M** (2010) Highly specific capture and direct MALDI MS analysis of phosphopeptides by zirconium phosphonate on self-assembled monolayers. *Anal Chem* 82:219-228

Horneffer V, Gluckmann M, Kruger B, **Karas M**, Strupat K, Hillenkamp F (2006) Matrix-analyte-interaction in MALDI-MS: Pellet and nano-electrospray preparations. *Int J Mass Spectrom* 249:426-432

Jaskolla T, Fuchs B, **Karas M**, Schiller J (2009) The new matrix 4-chloro-alpha-cyanocinnamic acid allows the detection of phosphatidylethanolamine chloramines by MALDI-TOF mass spectrometry. *J Am Soc Mass Spectr* 20:867-874

Jaskolla TW, **Karas M**, Roth U, Steinert K, Menzel C, Reihls K (2009) Comparison between vacuum sublimed matrices and conventional dried droplet preparation in MALDI-TOF mass spectrometry. *J Am Soc Mass Spectr* 20:1104-1114

Jaskolla TW, Lehmann WD, **Karas M** (2008) 4-Chloro- α -cyanocinnamic acid is an advanced, rationally designed MALDI matrix. *P Natl Acad Sci USA* 105:12200-12205

Jaskolla TW, Papasotiriou DG, **Karas M** (2009) Comparison between the Matrices alpha-Cyano-4-hydroxycinnamic Acid and 4-Chloro-alpha-cyanocinnamic Acid for Trypsin, Chymotrypsin, and Pepsin Digestions by MALDI-TOF Mass Spectrometry. *J Proteome Res* 8:3588-3597

Karas M (2007) Analytic - On the revolution on ripeness Nachr Chem, pp 1016-1019 vol 55)

Knauer O, Binai NA, Carra G, Beckhaus T, Hanschmann KM, Renne T, Backert S, **Karas M**, Wessler S (2008) Differential phosphoproteome profiling reveals a functional role for VASP in Helicobacter pylori-induced cytoskeleton turnover in gastric epithelial cells. *Cell Microbiol* 10:2285-2296

Kouyianou K, Aivaliotis M, Gevaert K, **Karas M**, Tsiotis G (2010) Membrane proteome of the green sulfur bacterium Chlorobium tepidum (syn. Chlorobaculum tepidum) analyzed by gel-based and gel-free methods. *Photosynth Res* 104:153-162

Kruger P, Daneshfar R, Eckert GP, Klein J, Volmer DA, Bahr U, Muller WE, **Karas M**, Schubert-Zsilavecz M, Abdel-Tawab M (2008) Metabolism of boswellic acids in vitro and in vivo. *Drug Metab Dispos* 36:1135-1142

- Ladig R, Sommer MS, Hahn A, Leisegang MS, Papatotiriou DG, Ibrahim M, Elkehal R, **Karas M**, Zickermann V, Gutensohn M, Brandt U, Klösigen RB, Schleiff E (2011) A high-definition native polyacrylamide gel electrophoresis system for the analysis of membrane complexes. *Plant J* 67:181-194
- Lehner I, Basting D, Meyer B, Haase W, Manolikas T, Kaiser C, **Karas M**, Glaubitz C (2008) The key residue for substrate transport (Glu(14)) in the EmrE dimer is asymmetric. *J Biol Chem* 283:3281-3288
- Meurer S, Pioch S, Pabst T, Opitz N, Schmidt PM, Beckhaus T, Wagner K, Matt S, Gegenbauer K, Geschka S, **Karas M**, Stasch JP, Schmidt H, Müller-Esterl W (2009) Nitric oxide-independent vasodilator rescues heme-oxidized soluble guanylate cyclase from proteasomal degradation. *Circ Res* 105:33-U91
- Meyer B, Wittig I, Trifilieff E, **Karas M**, Schägger H (2007) Identification of two proteins associated with mammalian ATP synthase. *Mol Cell Proteomics* 6:1690-1699
- Meyer B, Wurm JP, Kotter P, Leisegang MS, Schilling V, Buchhaupt M, Held M, Bahr U, **Karas M**, Heckel A, Bohnsack MT, Wöhnert J, Entian KD (2011) The Bowen-Conradi syndrome protein Nep1 (Emg1) has a dual role in eukaryotic ribosome biogenesis, as an essential assembly factor and in the methylation of Psi 1191 in yeast 18S rRNA. *Nucleic Acids Res* 39:1526-1537
- Morciano M, Beckhaus T, **Karas M**, Zimmermann H, Volkandt W (2009) The proteome of the presynaptic active zone: from docked synaptic vesicles to adhesion molecules and maxi-channels. *J Neurochem* 108:662-675
- Moritz A, Scheschonka A, Beckhaus T, **Karas M**, Betz H (2009) Metabotropic glutamate receptor 4 interacts with microtubule-associated protein 1B. *Biochem Biophys Res Commun* 390:82-86
- Paarmann I, Schmitt B, Meyer B, **Karas M**, Betz H (2006) Mass spectrometric analysis of glycine receptor-associated gephyrin splice variants. *J Biol Chem* 281:34918-34925
- Papatotiriou DG, Jaskolla TW, Markoutsas S, Baeumlisberger D, **Karas M**, Meyer B (2010) Peptide mass fingerprinting after less specific in-gel proteolysis using MALDI-LTQ-Orbitrap and 4-chloro-alpha-cyanocinnamic acid. *J Proteome Res* 9:2619-2629
- Papatotiriou DG, Markoutsas S, Meyer B, Papadioti A, **Karas M**, Tsiotis G (2008) Comparison of the membrane subproteomes during growth of a new Pseudomonas strain on lysogeny broth medium, glucose, and phenol. *J Proteome Res* 7:4278-4288
- Peng GH, Bostina M, Radermacher M, Rais I, **Karas M**, Michel H (2006) Biochemical and electron microscopic characterization of the F1F0 ATP Synthase from the hyperthermophilic eubacterium Aquifex aeolicus. *FEBS Lett* 580:5934-5940
- Persike M, **Karas M** (2009) Rapid simultaneous quantitative determination of different small pharmaceutical drugs using a conventional matrix-assisted laser desorption/ionization time-of-flight mass spectrometry system. *Rapid Commun Mass Sp* 23:3555-3562
- Persike M, Zimmermann M, Klein J, **Karas M** (2010) Quantitative determination of acetylcholine and choline in microdialysis samples by MALDI-TOF MS. *Anal Chem* 82:922-929
- Pfenninger A, Chan SY, **Karas M**, Finke B, Stahl B, Costello CE (2008) Mass spectrometric detection of multiple extended series of neutral highly fucosylated N-acetyllactosamine oligosaccharides in human milk. *Int J Mass Spectrom* 278:129-136
- Rietschel B, Arrey TN, Meyer B, Bornemann S, Schuerken M, **Karas M**, Poetsch A (2009) Elastase digests new ammunition for shotgun membrane proteomics. *Mol Cell Proteomics* 8:1029-1043
- Rietschel B, Baeumlisberger D, Arrey TN, Bornemann S, Rohmer M, Schuerken M, **Karas M**, Meyer B (2009) The benefit of combining nLC-MALDI-Orbitrap MS data with nLC-MALDI-TOF/TOF data for proteomic analyses employing elastase. *J Proteome Res* 8:5317-24

- Rietschel B, Bornemann S, Arrey TN, Baeumlisberger D, **Karas M**, Meyer B (2009) Membrane protein analysis using an improved peptic in-solution digestion protocol. *Proteomics* 9:5553-5557
- Rohmer M, Meyer B, Mank M, Stahl B, Bahr U, **Karas M** (2010) 3-Aminoquinoline acting as matrix and derivatizing agent for MALDI MS analysis of oligosaccharides. *Anal Chem* 82:3719-3726
- Rosenow A, Arrey TN, Bouwman FG, Noben JP, Wabitsch M, Mariman ECM, **Karas M**, Renes J (2010) Identification of novel human adipocyte secreted proteins by using SGBS Cells. *J Proteome Res* 9:5389-5401
- Schulz E, **Karas M**, Rosu F, Gabelica V (2006) Influence of the matrix on analyte fragmentation in atmospheric pressure MALDI. *J Am Soc Mass Spectr* 17:1005-1013
- Siegel C, Hallstrom T, Skerka C, Eberhardt H, Uzonyi B, Beckhaus T, **Karas M**, Wallich R, Stevenson B, Zipfel PF, Kraiczy P (2010) Complement Factor H-Related Proteins CFHR2 and CFHR5 Represent Novel Ligands for the Infection-Associated CRASP Proteins of *Borrelia burgdorferi*. *PLoS One* 5:14
- Strey CW, Gestrich J, Beckhaus T, Marquez-Pinilla RM, Oppermann E, Monch C, Lambris JD, **Karas M**, Bechstein WO (2010) Hypoxia and reoxygenation of primary human hepatocytes induce proteome changes of glucose metabolism, oxidative protection and peroxisomal function. *Int J Mol Med* 26:577-584
- Suhartono M, Weidlich M, Stein T, **Karas M**, Durner G, Gobel MW (2008) Synthesis of non-natural aromatic alpha-amino acids by a Heck reaction. *Eur J Org Chem*:1608-1614
- Tanrikulu Y, Nietert M, Scheffer U, Proschak E, Grabowski K, Schneider P, Weidlich M, **Karas M**, Gobel M, Schneider G (2007) Scaffold hopping by "Fuzzy" pharmacophores and its application to RNA targets. *ChemBioChem* 8:1932-+
- Tausch L, Henkel A, Siemoneit U, Poeckel D, Kather N, Franke L, Hofmann B, Schneider G, Angioni C, Geisslinger G, Skarke C, Holtmeier W, Beckhaus T, **Karas M**, Jauch J, Werz O (2009) Identification of human Cathepsin G as a functional target of boswellic acids from the anti-inflammatory remedy frankincense. *J Immunol* 183:3433-3442
- Teuber K, Schiller J, Fuchs B, **Karas M**, Jaskolla TW (2010) Significant sensitivity improvements by matrix optimization: a MALDI-TOF mass spectrometric study of lipids from hen egg yolk. *Chem Phys Lipids* 163:552-560
- Tsirogiani E, Aivaliotis M, Papisotiriou DG, **Karas M**, Tsiotis G (2006) Identification of inducible protein complexes in the phenol degrader *Pseudomonas* sp strain phDV1 by blue native gel electrophoresis and mass spectrometry. *Amino Acids* 30:63-72
- Urbanek DA, Proschak E, Tanrikulu Y, Becker S, **Karas M**, Schneider G (2011) Scaffold-hopping from aminoglycosides to small synthetic inhibitors of bacterial protein biosynthesis using a pseudoreceptor model. *MedChemComm* 2:181-184
- Vettermann C, Castor D, Mekker A, Gerrits B, **Karas M**, Jack HM (2011) Proteome profiling suggests a pro-inflammatory role for plasma cells through release of high-mobility group box 1 protein. *Proteomics* 11:1228-1237
- Weidlich M, Klammt C, Bernhard F, **Karas M**, Stein T (2008) Differential activity profiles of translation inhibitors in whole-cell and cell-free approaches. *Letts Appl Microbiol* 46:155-159
- Wittig I, Beckhaus T, Wumaier Z, **Karas M**, Schagger H (2010) Mass estimation of native proteins by blue native electrophoresis principles and practical hints. *Mol Cell Proteomics* 9:2149-2161
- Wittig I, **Karas M**, Schagger H (2007) High resolution clear native electrophoresis for in-gel functional assays and fluorescence studies of membrane protein complexes. *Mol Cell Proteomics* 6:1215-1225
- Wohlgemuth J, **Karas M**, Eichhorn T, Hendriks R, Andrecht S (2009) Quantitative site-specific analysis of protein glycosylation by LC-MS using different glycopeptide-enrichment strategies. *Anal Biochem* 395:178-188

Wohlgemuth J, **Karas M**, Jiang W, Hendriks R, Andrecht S (2010) Enhanced glyco-profiling by specific glycopeptide enrichment and complementary monolithic nano-LC (ZIC-HILIC/RP18e)/ESI-MS analysis. *J Sep Sci* 33:880-890

Wurm JP, Meyer B, Bahr U, Held M, Frolow O, Kotter P, Engels JW, Heckel A, **Karas M**, Entian KD, Wöhnert J (2010) The ribosome assembly factor Nep1 responsible for Bowen-Conradi syndrome is a pseudouridine-N1-specific methyltransferase. *Nucleic Acids Res* 38:2387-2398